



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 9/02 // (C12N 9/02, C12R 1:645)	A1	(11) International Publication Number: WO 98/27197 (43) International Publication Date: 25 June 1998 (25.06.98)
(21) International Application Number: PCT/DK97/00566 (22) International Filing Date: 12 December 1997 (12.12.97) (30) Priority Data: 1450/96 19 December 1996 (19.12.96) DK 1020/97 8 September 1997 (08.09.97) DK (71) Applicant: NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK). (72) Inventors: PEDERSEN, Anders, Hjelholt; Nybro Vænge 58, DK-2800 Lyngby (DK). SVENDSEN, Allan; Bakkeledet 28, DK-3460 Birkerød (DK). SCHNEIDER, Palle; Rydtoften 43, DK-2750 Bellerup (DK). RASMUSSEN, Grethe; Brudedalen 1, DK-3520 Farum (DK). CHERRY, Joel, R.; C.E.V. Knuthsvej 9, DK-2100 Hellerup (DK).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: MYCELIOPHTHORA AND SCYTALIDIUM LACCASE VARIANTS HAVING IMPROVED STABILITY		
(57) Abstract The present invention relates to laccase mutants with improved stability properties, in particular to <i>Myceliophthora</i> and <i>Scytalidium</i> laccase variants comprising mutations in one or more tyrosine, tryptophan or methionine residues.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KP	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire		Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

MYCELIOPHTHORA AND SCYTALIDIUM LACCASE VARIANTS HAVING IMPROVED STABILITY**FIELD OF THE INVENTION**

5 The present invention relates to laccase mutants with improved stability properties.

BACKGROUND OF THE INVENTION

10 Laccase is a polyphenol oxidase (EC 1.10.3.2) which catalyses the oxidation of a variety of inorganic and aromatic compounds, particularly phenols, with the concomitant reduction of molecular oxygen to water.

Laccase belongs to a family of blue copper-containing
15 oxidases which includes ascorbate oxidase and the mammalian plasma protein ceruloplasmin. All these enzymes are multi-copper-containing proteins.

Because laccases are able to catalyze the oxidation of a variety of inorganic and aromatic compounds, laccases have been
20 suggested in many potential industrial applications such as lignin modification, paper strengthening, dye transfer inhibition in detergents, phenol polymerization, hair colouring, and waste water treatment. A major problem with the use of laccases are their poor storage stability at temperatures above room
25 temperature, especially at 40°C.

In Example 1 of the present application we have tested the stability of *Myceliophthora thermophila* laccase at 40°C, and it can be seen that after 2 weeks of storage the laccase activity is down to less than 50% of the initial value. For many purposes
30 such a decrease is unacceptable, so it is the purpose of the present invention to create laccase variants with improved stability.

BRIEF DISCLOSURE OF THE INVENTION

35 The present invention relates to laccase variants, in particular to a variant of a parent laccase, which variant has laccase activity, improved stability as compared to said parent laccase,

and comprises a mutation in one or more tyrosine, tryptophan or methionine residues, wherein the parent laccase has the amino acid sequence given in SEQ ID No. 1 or the parent laccase has an amino acid sequence which is at least 80% homologous to SEQ ID No. 1.

In still further aspects the invention relates to DNA encoding such variants and methods of preparing the variants.

Finally, the invention relates to the use of the variants for various industrial purposes.

DETAILED DISCLOSURE OF THE INVENTION

Homologous Laccases

A number of laccases produced by different fungi are homologous on the amino acid level. For instance, when using the homology percent obtained from UWGCG program using the GAP program with the default parameters (penalties: gap weight=3.0, length weight=0.1; WISCONSIN PACKAGE Version 8.1-UNIX, August 1995, Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA 53711) the following homology was found:

Myceliophthora thermophila laccase comprising the amino acid sequence shown in SEQ ID No. 1: 100%;

Scytalidium thermophilum laccase comprising the amino acid sequence shown in SEQ ID No. 2: 81.2%.

Because of the homology found between the above mentioned laccases, they are considered to belong to the same class of laccases, namely the class of "Myceliophthora-like laccases".

Accordingly, in the present context, the term "Myceliophthora-like laccase" is intended to indicate a laccase which, on the amino acid level, displays a homology of at least 80% to the *Myceliophthora* laccase SEQ ID NO 1, or a laccase which, on the amino acid level, displays a homology of at least 85% to the *Myceliophthora* laccase SEQ ID NO 1, or a laccase which, on the amino acid level, displays a homology of at least 90% to the *Myceliophthora* laccase SEQ ID NO 1, or a laccase

which, on the amino acid level, displays a homology of at least 95% to the *Myceliophthora* laccase SEQ ID NO 1, or a laccase which, on the amino acid level, displays a homology of at least 98% to the *Myceliophthora* laccase SEQ ID NO 1.

5

In the present context, "derived from" is intended not only to indicate a laccase produced or producible by a strain of the organism in question, but also a laccase encoded by a DNA sequence isolated from such strain and produced in a host organism containing said DNA sequence. Finally, the term is intended to indicate a laccase which is encoded by a DNA sequence of synthetic and/or cDNA origin and which has the identifying characteristics of the laccase in question.

15 Variants with altered stability

It is contemplated that it is possible to improve the stability of a parent *Myceliophthora* laccase or a parent *Myceliophthora*-like laccase by making variants:

Such a variant has laccase activity, improved stability as compared to said parent laccase, and comprises a mutation in one or more tyrosine, tryptophan or methionine residues. The parent laccase has the amino acid sequence given in SEQ ID No. 1 or the parent laccase has an amino acid sequence which is at least 80% homologous to SEQ ID No. 1.

25 Preferred positions for mutations in *Myceliophthora thermophila* laccase (SEQ ID No 1) and in *Scytalidium thermophilum* laccase (SEQ ID No 2) are the following:

Myceliophthora thermophila:

30 M433,
W373,
W136,
Y145,
M480,
35 Y137,
Y176,
M254, and/or

W507.

Scytalidium thermophilum:

M483,
5 W422,
W181,
Y190,
M530,
Y182,
10 Y221,
M300, and/or
M313.

In particular the following mutations in *Myceliophthora*
15 *thermophila* laccase (SEQ ID No 1) and in *Scytalidium thermophilum*
laccase (SEQ ID No 2) are preferred:

A variant of a parent *Myceliophthora thermophila* laccase,
which comprises a substitution in a position corresponding to at
least one of the following positions in SEQ ID No. 1:

20 M433 A, V, L, I, P, F, W, G, S, T, C, Y, N, Q, D, E, K, R, H;
W373 A, V, L, I, P, F, M, G, S, T, C, Y, N, Q, D, E, K, R, H;
W136 A, V, L, I, P, F, M, G, S, T, C, Y, N, Q, D, E, K, R, H;
Y145 A, V, L, I, P, F, W, G, S, T, C, M, N, Q, D, E, K, R, H;
M480 A, V, L, I, P, F, W, G, S, T, C, Y, N, Q, D, E, K, R, H;
25 Y137 A, V, L, I, P, F, W, G, S, T, C, M, N, Q, D, E, K, R, H;
Y176 A, V, L, I, P, F, W, G, S, T, C, M, N, Q, D, E, K, R, H;
M254 A, V, L, I, P, F, W, G, S, T, C, Y, N, Q, D, E, K, R, H;
W507 A, V, L, I, P, F, M, G, S, T, C, Y, N, Q, D, E, K, R, H;

30 in particular at least one of the following positions in SEQ ID
No. 1:

M433 F, V, I, L, Q;
W373 F, H;
W136 F, H;
35 Y145 F;
M480 F, V, I, L, Q;
Y137 F;

Y176 F;
 M254 F, V, I, L, Q; and/or
 W507 F, H.

5 A variant of a parent *Scytalidium thermophilum* laccase, which comprises a substitution in a position corresponding to at least one of the following positions in SEQ ID No. 2:

M483 A, V, L, I, P, F, W, G, S, T, C, Y, N, Q, D, E, K, R, H;
 W422 A, V, L, I, P, F, M, G, S, T, C, Y, N, Q, D, E, K, R, H;
 10 W181 A, V, L, I, P, F, M, G, S, T, C, Y, N, Q, D, E, K, R, H;
 Y190 A, V, L, I, P, F, W, G, S, T, C, M, N, Q, D, E, K, R, H;
 M530 A, V, L, I, P, F, W, G, S, T, C, Y, N, Q, D, E, K, R, H;
 Y182 A, V, L, I, P, F, W, G, S, T, C, M, N, Q, D, E, K, R, H;
 Y221 A, V, L, I, P, F, W, G, S, T, C, M, N, Q, D, E, K, R, H;
 15 M300 A, V, L, I, P, F, W, G, S, T, C, Y, N, Q, D, E, K, R, H;
 M313 A, V, L, I, P, F, W, G, S, T, C, Y, N, Q, D, E, K, R, H;

in particular at least one of following positions in SEQ ID No. 2:

20 M483 F, V, I, L, Q;
 W422 F, H;
 W181 F, H;
 Y190 F;
 M530 F, V, I, L, Q;
 25 Y182 F;
 Y221 F;
 M300 F, V, I, L, Q; and/or
 M313 F, V, I, L, Q.

30 Methods of preparing laccase variants

Several methods for introducing mutations into genes are known in the art. After a brief discussion of the cloning of laccase-encoding DNA sequences, methods for generating mutations at specific sites within the laccase-encoding sequence will be
 35 discussed.

Cloning a DNA sequence encoding a laccase

The DNA sequence encoding a parent laccase may be isolated from any cell or microorganism producing the laccase in question, using various methods well known in the art. First, a genomic DNA and/or cDNA library should be constructed using chromosomal DNA or messenger RNA from the organism that produces the laccase to be studied. Then, if the amino acid sequence of the laccase is known, homologous, labelled oligonucleotide probes may be synthesized and used to identify laccase-encoding clones from a genomic library prepared from the organism in question. Alternatively, a labelled oligonucleotide probe containing sequences homologous to a known laccase gene could be used as a probe to identify laccase-encoding clones, using hybridization and washing conditions of lower stringency.

A method for identifying laccase-encoding clones involves inserting cDNA into an expression vector, such as a plasmid, transforming laccase-negative fungi with the resulting cDNA library, and then plating the transformed fungi onto agar containing a substrate for laccase, thereby allowing clones expressing the laccase to be identified.

Alternatively, the DNA sequence encoding the enzyme may be prepared synthetically by established standard methods, e.g. the phosphoroamidite method. In the phosphoroamidite method, oligonucleotides are synthesized, e.g. in an automatic DNA synthesizer, purified, annealed, ligated and cloned in appropriate vectors.

Finally, the DNA sequence may be of mixed genomic and synthetic origin, mixed synthetic and cDNA origin or mixed genomic and cDNA origin, prepared by ligating fragments of synthetic, genomic or cDNA origin (as appropriate, the fragments corresponding to various parts of the entire DNA sequence), in accordance with standard techniques. The DNA sequence may also be prepared by polymerase chain reaction (PCR) using specific primers.

Site-directed mutagenesis

Once a laccase-encoding DNA sequence has been isolated, and desirable sites for mutation identified, mutations may be introduced using synthetic oligonucleotides. These oligonucleotides contain nucleotide sequences flanking the desired mutation sites; mutant nucleotides are inserted during oligonucleotide synthesis. In a specific method, a single-stranded gap of DNA, bridging the laccase-encoding sequence, is created in a vector carrying the laccase gene. Then the synthetic nucleotide, bearing the desired mutation, is annealed to a homologous portion of the single-stranded DNA. The remaining gap is then filled in with T7 DNA polymerase and the construct is ligated using T4 ligase. A specific example of this method is described in Morinaga et al. (1984). US 4,760,025 discloses the introduction of oligonucleotides encoding multiple mutations by performing minor alterations of the cassette. However, an even greater variety of mutations can be introduced at any one time by the Morinaga method, because a multitude of oligonucleotides, of various lengths, can be introduced.

Another method of introducing mutations into laccase-encoding DNA sequences is described in Nelson and Long (1989). It involves the 3-step generation of a PCR fragment containing the desired mutation introduced by using a chemically synthesized DNA strand as one of the primers in the PCR reactions. From the PCR-generated fragment, a DNA fragment carrying the mutation may be isolated by cleavage with restriction endonucleases and reinserted into an expression plasmid.

Random mutagenesis

The random mutagenesis of a DNA sequence encoding a parent laccase may conveniently be performed by use of any method known in the art.

For instance, the random mutagenesis may be performed by use of a suitable physical or chemical mutagenizing agent, by use of a suitable oligonucleotide, or by subjecting the DNA sequence to PCR generated mutagenesis. Furthermore, the random mutagenesis may be performed by use of any combination of these mutagenizing agents.

The mutagenizing agent may, e.g., be one which induces transitions, transversions, inversions, scrambling, deletions, and/or insertions.

Examples of a physical or chemical mutagenizing agent suitable for the present purpose include ultraviolet (UV) irradiation, hydroxylamine, N-methyl-N'-nitro-N-nitrosoguanidine (MNNG), O-methyl hydroxylamine, nitrous acid, ethyl methane sulphonate (EMS), sodium bisulphite, formic acid, and nucleotide analogues.

10 When such agents are used, the mutagenesis is typically performed by incubating the DNA sequence encoding the parent enzyme to be mutagenized in the presence of the mutagenizing agent of choice under suitable conditions for the mutagenesis to take place, and selecting for mutated DNA having the desired
15 properties.

When the mutagenesis is performed by the use of an oligonucleotide, the oligonucleotide may be doped or spiked with the three non-parent nucleotides during the synthesis of the oligonucleotide at the positions which are to be changed. The
20 doping or spiking may be done so that codons for unwanted amino acids are avoided. The doped or spiked oligonucleotide can be incorporated into the DNA encoding the laccase enzyme by any published technique, using e.g. PCR, LCR or any DNA polymerase and ligase.

25 When PCR-generated mutagenesis is used, either a chemically treated or non-treated gene encoding a parent laccase enzyme is subjected to PCR under conditions that increase the misincorporation of nucleotides (Deshler 1992; Leung et al., Technique, Vol.1, 1989, pp. 11-15).

30 A mutator strain of *E. coli* (Fowler et al., Molec. Gen. Genet., 133, 1974, pp. 179-191), *S. cerevisiae* or any other microbial organism may be used for the random mutagenesis of the DNA encoding the laccase enzyme by e.g. transforming a plasmid containing the parent enzyme into the mutator strain, growing the
35 mutator strain with the plasmid and isolating the mutated plasmid from the mutator strain. The mutated plasmid may subsequently be transformed into the expression organism.

The DNA sequence to be mutagenized may conveniently be present in a genomic or cDNA library prepared from an organism expressing the parent laccase enzyme. Alternatively, the DNA sequence may be present on a suitable vector such as a plasmid or a bacteriophage, which as such may be incubated with or otherwise exposed to the mutagenizing agent. The DNA to be mutagenized may also be present in a host cell either by being integrated in the genome of said cell or by being present on a vector harboured in the cell. Finally, the DNA to be mutagenized may be in isolated form. It will be understood that the DNA sequence to be subjected to random mutagenesis is preferably a cDNA or a genomic DNA sequence.

In some cases it may be convenient to amplify the mutated DNA sequence prior to the expression step or the screening step being performed. Such amplification may be performed in accordance with methods known in the art, the presently preferred method being PCR-generated amplification using oligonucleotide primers prepared on the basis of the DNA or amino acid sequence of the parent enzyme.

Subsequent to the incubation with or exposure to the mutagenizing agent, the mutated DNA is expressed by culturing a suitable host cell carrying the DNA sequence under conditions allowing expression to take place. The host cell used for this purpose may be one which has been transformed with the mutated DNA sequence, optionally present on a vector, or one which was carried the DNA sequence encoding the parent enzyme during the mutagenesis treatment. Examples of suitable host cells are fungal hosts such as *Aspergillus niger* or *Aspergillus oryzae*.

The mutated DNA sequence may further comprise a DNA sequence encoding functions permitting expression of the mutated DNA sequence.

Localized random mutagenesis

The random mutagenesis may advantageously be localized to a part of the parent laccase in question. This may, e.g., be advantageous when certain regions of the enzyme have been identified to be of particular importance for a given property of

the enzyme, and when modified are expected to result in a variant having improved properties. Such regions may normally be identified when the tertiary structure of the parent enzyme has been elucidated and related to the function of the enzyme.

5 The localized random mutagenesis is conveniently performed by use of PCR-generated mutagenesis techniques as described above or any other suitable technique known in the art.

Alternatively, the DNA sequence encoding the part of the DNA sequence to be modified may be isolated, e.g. by being inserted
10 into a suitable vector, and said part may subsequently be subjected to mutagenesis by use of any of the mutagenesis methods discussed above.

With respect to the screening step in the above-mentioned method of the invention, this may conveniently be performed by
15 use of aa filter assay based on the following principle:

A microorganism capable of expressing the mutated laccase enzyme of interest is incubated on a suitable medium and under suitable conditions for the enzyme to be secreted, the medium being provided with a double filter comprising a first protein-
20 binding filter and on top of that a second filter exhibiting a low protein binding capability. The microorganism is located on the second filter. Subsequent to the incubation, the first filter comprising enzymes secreted from the microorganisms is separated from the second filter comprising the microorganisms. The first
25 filter is subjected to screening for the desired enzymatic activity and the corresponding microbial colonies present on the second filter are identified.

The filter used for binding the enzymatic activity may be any protein binding filter e.g. nylon or nitrocellulose. The top
30 filter carrying the colonies of the expression organism may be any filter that has no or low affinity for binding proteins e.g. cellulose acetate or Durapore™. The filter may be pretreated with any of the conditions to be used for screening or may be treated during the detection of enzymatic activity.

35 The enzymatic activity may be detected by a dye, fluorescence, precipitation, pH indicator, IR-absorbance or any other known technique for detection of enzymatic activity.

The detecting compound may be immobilized by any immobilizing

agent, e.g., agarose, agar, gelatine, polyacrylamide, starch, filter paper, cloth; or any combination of immobilizing agents.

Testing of variants of the invention

5 The storage stability of *Myceliophthora* variants or *Myceliophthora*-like variants should be investigated at 40°C for 2 weeks at pH 5, 8 and 9.3, respectively. The stability of the parent laccase and the variants may be tested both in a liquid buffer formulation and in a lyophilized form.

10 According to the invention the residual activity of the variants following two weeks of incubation are then compared to the residual activity of the parent laccase, and variants with an improved stability at either pH 5, 8 or 9.3 are selected.

15 Laccase activity

In the context of this invention, the laccase activity was measured using 10-(2-hydroxyethyl)-phenoxazine (HEPO) as substrate for the various laccases. HEPO was synthesized using the same procedure as described for 10-(2-hydroxyethyl)-
20 phenothiazine, (G. Cauquil in Bulletin de la Society Chimique de France, 1960, p. 1049). In the presence of oxygen laccases (E.C. 1.10.3.2) oxidize HEPO to a HEPO radical that can be monitored photometrically at 528 nm.

The *Myceliophthora thermophila* laccase was measured using
25 0.4 mM HEPO in 25 mM Tris-HCl, pH 7.5, 0.05% TWEEN-20 at 30 °C. The absorbance at 528 nm was followed for 200 s and the rate calculated from the linear part of the progress curve.

30 Expression of laccase variants

According to the invention, a DNA sequence encoding the variant produced by methods described above, or by any alternative methods known in the art, can be expressed, in enzyme form, using an expression vector which typically includes control
35 sequences encoding a promoter, operator, ribosome binding site, translation initiation signal, and, optionally, a repressor gene or various activator genes.

The recombinant expression vector carrying the DNA sequence encoding a laccase variant of the invention may be any vector which may conveniently be subjected to recombinant DNA procedures, and the choice of vector will often depend on the host cell into which it is to be introduced. Thus, the vector may be an autonomously replicating vector, i.e. a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid, a bacteriophage or an extrachromosomal element, minichromosome or an artificial chromosome. Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated.

In the vector, the DNA sequence should be operably connected to a suitable promoter sequence. The promoter may be any DNA sequence which shows transcriptional activity in the host cell of choice and may be derived from genes encoding proteins either homologous or heterologous to the host cell. Examples of suitable promoters for directing the transcription of the DNA sequence encoding a laccase variant of the invention, especially in a fungal host, are those derived from the gene encoding *A. oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *A. niger* neutral α -amylase, *A. niger* acid stable α -amylase, *A. niger* glucoamylase, *Rhizomucor miehei* lipase, *A. oryzae* alkaline protease, *A. oryzae* triose phosphate isomerase or *A. nidulans* acetamidase.

The expression vector of the invention may also comprise a suitable transcription terminator and, in eukaryotes, polyadenylation sequences operably connected to the DNA sequence encoding the laccase variant of the invention. Termination and polyadenylation sequences may suitably be derived from the same sources as the promoter.

The vector may further comprise a DNA sequence enabling the vector to replicate in the host cell in question. Examples of such sequences are the origins of replication of plasmids pUC19, pACYC177, pUB110, pE194, pAMB1 and pIJ702.

The vector may also comprise a selectable marker, e.g. a gene, the product of which complements a defect in the host cell,

such as one which confers antibiotic resistance such as ampicillin, kanamycin, chloramphenicol or tetracycline resistance. Furthermore, the vector may comprise *Aspergillus* selection markers such as amdS, argB, niaD and sC, a marker giving rise to 5 hygromycin resistance, or the selection may be accomplished by co-transformation, e.g. as described in WO 91/17243.

The procedures used to ligate the DNA construct of the invention encoding a laccase variant, the promoter, terminator and other elements, respectively, and to insert them into suitable 10 vectors containing the information necessary for replication, are well known to persons skilled in the art (cf., for instance, Sambrook et al. (1989)).

The cell of the invention, either comprising a DNA construct or an expression vector of the invention as defined above, is 15 advantageously used as a host cell in the recombinant production of a laccase variant of the invention. The cell may be transformed with the DNA construct of the invention encoding the variant, conveniently by integrating the DNA construct (in one or more copies) in the host chromosome. This integration is 20 generally considered to be an advantage as the DNA sequence is more likely to be stably maintained in the cell. Integration of the DNA constructs into the host chromosome may be performed according to conventional methods, e.g. by homologous or heterologous recombination. Alternatively, the cell may be 25 transformed with an expression vector as described above in connection with the different types of host cells.

The cell of the invention may be a cell of a higher organism such as a mammal or an insect, but is preferably a microbial cell, e.g. a fungal cell.

30 The filamentous fungus may advantageously belong to a species of *Aspergillus*, e.g. *Aspergillus oryzae* or *Aspergillus niger*. Fungal cells may be transformed by a process involving protoplast formation and transformation of the protoplasts followed by regeneration of the cell wall in a manner known per se. A 35 suitable procedure for transformation of *Aspergillus* host cells is described in EP 238 023.

In a yet further aspect, the present invention relates to a

method of producing a laccase variant of the invention, which method comprises cultivating a host cell as described above under conditions conducive to the production of the variant and recovering the variant from the cells and/or culture medium.

5 The medium used to cultivate the cells may be any conventional medium suitable for growing the host cell in question and obtaining expression of the laccase variant of the invention. Suitable media are available from commercial suppliers or may be prepared according to published recipes (e.g. as described in
10 catalogues of the American Type Culture Collection).

The laccase variant secreted from the host cells may conveniently be recovered from the culture medium by well-known procedures, including separating the cells from the medium by centrifugation or filtration, and precipitating proteinaceous
15 components of the medium by means of a salt such as ammonium sulphate, followed by the use of chromatographic procedures such as ion exchange chromatography, affinity chromatography, or the like.

20 Industrial Applications

The laccase variants of this invention possesses valuable properties allowing for various industrial applications, in particular lignin modification, paper strengthening, dye transfer inhibition in detergents, phenol polymerization, hair dyeing,
25 bleaching of textiles (in particular bleaching of denim as described in WO 96/12845 and WO 96/12846) and waste water treatment. Any detergent composition normally used for enzymes may be used, e.g., the detergent compositions disclosed in WO 95/01426.

30

The invention is further illustrated in the following examples, which are not intended to be in any way limiting to the scope of the invention as claimed.

35 EXAMPLE 1

Storage stability of the *Myceliophthora thermophila* laccase

The storage stability of the *Myceliophthora thermophila*

laccase was tested for 2 weeks at 40°C at pH 5, 8 and 9.3, respectively. The laccase (1 mg/ml) was dialyzed against 0.1 M sodium acetate, pH 5, or 0.1 M Tris-maleate, pH 8, or 0.1 M Tris-maleate, pH 9.3. Following dialysis the different preparations were poured into two sets of glass vials with screw caps: one for the liquid formulation and the other one for the lyophilized form. After two weeks of incubation the enzyme activity was measured as described above and the residual activity of the enzyme was calculated in percentage using a preparation of *Myceliophthora thermophila* kept at 4°C as a reference.

Table 1 Storage stability of *Myceliophthora thermophila*

pH	Liquid formulation		Lyophilized form	
	Residual activity (%)		Residual activity (%)	
5.0	<5		<5	
8.0	<5		<5	
9.3	35		30	

15

EXAMPLE 2

Storage stability of *Myceliophthora thermophila* variants

20 Laccase activity:

In this Example the *Myceliophthora thermophila* laccase variants were measured using 0.4 mM HEPO in 0.1 M Tris-maleate, pH 7.5, 0.05% TWEEN-20 at 30°C. The absorbance at 528 nm was followed for 200 s and the rate calculated from the linear part of the progress curve.

The storage stability of the *Myceliophthora thermophila* variants were tested for 4 weeks at 40°C at pH 5, 7, and 9.3, respectively. The laccase (1 mg/ml) was dialyzed against 0.1 M Tris-maleate, pH 5 or 0.1 M Tris-maleate, pH 7 or 0.1 M Tris-maleate, pH 9.3. Following dialysis the different preparations were poured into two set of glass vials with screw caps: one

for the liquid formulation and the other set of glasses for lyophilization. Following two and four weeks of incubation the enzyme activity was measured as described above and the residual activity of the variants were calculated in percentage using a preparation kept at 4°C as reference.

Table 2. Storage stability of *Myceliophthora thermophila* variants, lyophilized formulation

	Residual activity, pH 5		Residual activity, pH 7		Residual activity, pH 9.2	
	2 weeks	4 weeks	2 weeks	4 weeks	2 weeks	4 weeks
wt	18	18	55	36	59	38
W136F	<5	<5	76	64	88	77
Y137F	12	<5	58	41	64	49
Y145F	<5	<5	53	20	45	51
W373F	14	14	33	19	51	36
M433I	7	<5	57	43	74	35
M480L	33	18	65	32	72	52
W507F	18	<5	72	51	68	71

10

In lyophilized form none of the tested variants have improved stability at pH 5. At pH 7 and pH 9.2 both W136F and W507F have increased stability. At pH 9.2 M480L is also better than wt.

15

Table 3. Storage stability of *Myceliophthora thermophila* variants, liquid formulation

	Residual activity, pH 5, 2 weeks	Residual activity, pH 7, 2 weeks	Residual activity, pH 9.2, 2 weeks
wt	<5	5	20
W136F	5	28	55

Y137F	<5	<5	<5
Y145F	<5	<5	<5
W373F	<5	40	<5
M433I	8	40	65
M480L	<5	<5	15
W507F	<5	<5	22

Also in the liquid formulation none of the tested variants have improved stability at pH 5. At pH 7 and pH 9.2 both
5 W136F and M433I has increased stability. At pH7 W373F has better stability than wt but the variant loses the stability completely at pH 9.2.

Of the tested variants only W136F has increased stability in
10 both formulations.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT:

- (A) NAME: NOVO NORDISK A/S
(B) STREET: Novo Alle
(C) CITY: Bagsvaerd
(E) COUNTRY: Denmark
(F) POSTAL CODE (ZIP): DK-2880
(G) TELEPHONE: +45 44 44 88 88
(H) TELEFAX: +45 44 49 05 55

10

15 (ii) TITLE OF INVENTION: LACCASE MUTANTS

(iii) NUMBER OF SEQUENCES: 2

(iv) COMPUTER READABLE FORM:

20

- (A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

25

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

30

- (A) LENGTH: 573 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

40

Gln Gln Ser Cys Asn Thr Pro Ser Asn Arg Ala Cys Trp Thr Asp Gly
1 5 10 15

Tyr Asp Ile Asn Thr Asp Tyr Glu Val Asp Ser Pro Asp Thr Gly Val
20 25 30

45

	Val	Arg	Pro	Tyr	Thr	Leu	Thr	Leu	Thr	Glu	Val	Asp	Asn	Trp	Thr	Gly	
			35					40					45				
5	Pro	Asp	Gly	Val	Val	Lys	Glu	Lys	Val	Met	Leu	Val	Asn	Asn	Ser	Ile	
		50					55					60					
	Ile	Gly	Pro	Thr	Ile	Phe	Ala	Asp	Trp	Gly	Asp	Thr	Ile	Gln	Val	Thr	
	65					70					75				80		
10	Val	Ile	Asn	Asn	Leu	Glu	Thr	Asn	Gly	Thr	Ser	Ile	His	Trp	His	Gly	
					85					90					95		
	Leu	His	Gln	Lys	Gly	Thr	Asn	Leu	His	Asp	Gly	Ala	Asn	Gly	Ile	Thr	
15					100				105					110			
	Glu	Cys	Pro	Ile	Pro	Pro	Lys	Gly	Gly	Arg	Lys	Val	Tyr	Arg	Phe	Lys	
			115					120					125				
20	Ala	Gln	Gln	Tyr	Gly	Thr	Ser	Trp	Tyr	His	Ser	His	Phe	Ser	Ala	Gln	
			130					135					140				
	Tyr	Gly	Asn	Gly	Val	Val	Gly	Ala	Ile	Gln	Ile	Asn	Gly	Pro	Ala	Ser	
	145					150					155				160		
25	Leu	Pro	Tyr	Asp	Thr	Asp	Leu	Gly	Val	Phe	Pro	Ile	Ser	Asp	Tyr	Tyr	
						165				170					175		
	Tyr	Ser	Ser	Ala	Asp	Glu	Leu	Val	Glu	Leu	Thr	Lys	Asn	Ser	Gly	Ala	
30				180					185					190			
	Pro	Phe	Ser	Asp	Asn	Val	Leu	Phe	Asn	Gly	Thr	Ala	Lys	His	Pro	Glu	
			195					200					205				
35	Thr	Gly	Glu	Gly	Glu	Tyr	Ala	Asn	Val	Thr	Leu	Thr	Pro	Gly	Arg	Arg	
		210					215						220				
	His	Arg	Leu	Arg	Leu	Ile	Asn	Thr	Ser	Val	Glu	Asn	His	Phe	Gln	Val	
	225				230						235				240		
40	Ser	Leu	Val	Asn	His	Thr	Met	Cys	Ile	Ile	Ala	Ala	Asp	Met	Val	Pro	
					245					250					255		
	Val	Asn	Ala	Met	Thr	Val	Asp	Ser	Leu	Phe	Leu	Gly	Val	Gly	Gln	Arg	
45				260					265					270			

Tyr Asp Val Val Ile Glu Ala Asn Arg Thr Pro Gly Asn Tyr Trp Phe
 275 280 285

5 Asn Val Thr Phe Gly Gly Gly Leu Leu Cys Gly Gly Ser Arg Asn Pro
 290 295 300

Tyr Pro Ala Ala Ile Phe His Tyr Ala Gly Ala Pro Gly Gly Pro Pro
 305 310 315 320

10 Thr Asp Glu Gly Lys Ala Pro Val Asp His Asn Cys Leu Asp Leu Pro
 325 330 335

15 Asn Leu Lys Pro Val Val Ala Arg Asp Val Pro Leu Ser Gly Phe Ala
 340 345 350

Lys Arg Ala Asp Asn Thr Leu Asp Val Thr Leu Asp Thr Thr Gly Thr
 355 360 365

20 Pro Leu Phe Val Trp Lys Val Asn Gly Ser Ala Ile Asn Ile Asp Trp
 370 375 380

Gly Arg Ala Val Val Asp Tyr Val Leu Thr Gln Asn Thr Ser Phe Pro
 385 390 395 400

25 Pro Gly Tyr Asn Ile Val Glu Val Asn Gly Ala Asp Gln Trp Ser Tyr
 405 410 415

30 Trp Leu Ile Glu Asn Asp Pro Gly Ala Pro Phe Thr Leu Pro His Pro
 420 425 430

Met His Leu His Gly His Asp Phe Tyr Val Leu Gly Arg Ser Pro Asp
 435 440 445

35 Glu Ser Pro Ala Ser Asn Glu Arg His Val Phe Asp Pro Ala Arg Asp
 450 455 460

Ala Gly Leu Leu Ser Gly Ala Asn Pro Val Arg Arg Asp Val Ser Met
 465 470 475 480

40 Leu Pro Ala Phe Gly Trp Val Val Leu Ser Phe Arg Ala Asp Asn Pro
 485 490 495

45 Gly Ala Trp Leu Phe His Cys His Ile Ala Trp His Val Ser Gly Gly
 500 505 510

Leu Gly Val Val Tyr Leu Glu Arg Ala Asp Asp Leu Arg Gly Ala Val
 515 520 525

5 Ser Asp Ala Asp Ala Asp Asp Leu Asp Arg Leu Cys Ala Asp Trp Arg
 530 535 540

Arg Tyr Trp Pro Thr Asn Pro Tyr Pro Lys Ser Asp Ser Gly Leu Lys
 545 550 555 560

10

His Arg Trp Val Glu Glu Gly Glu Trp Leu Val Lys Ala
 565 570

(2) INFORMATION FOR SEQ ID NO: 2:

15

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 616 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

20

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

30 Met Lys Arg Phe Phe Ile Asn Ser Leu Leu Leu Leu Ala Gly Leu Leu
 1 5 10 15

Asn Ser Gly Ala Leu Ala Ala Pro Ser Thr His Pro Arg Ser Asn Pro
 20 25 30

35 Asp Ile Leu Leu Glu Arg Asp Asp His Ser Leu Thr Ser Arg Gln Gly
 35 40 45

Ser Cys His Ser Pro Ser Asn Arg Ala Cys Trp Cys Ser Gly Phe Asp
 50 55 60

40

Ile Asn Thr Asp Tyr Glu Thr Lys Thr Pro Asn Thr Gly Val Val Arg
 65 70 75 80

45 Arg Tyr Thr Phe Asp Ile Thr Glu Val Asp Asn Arg Pro Gly Pro Asp
 85 90 95

Gly Val Ile Lys Glu Lys Leu Met Leu Ile Asn Asp Lys Leu Leu Gly
 100 105 110

5 Pro Thr Val Phe Ala Asn Trp Gly Asp Thr Ile Glu Val Thr Val Asn
 115 120 125

Asn His Leu Arg Thr Asn Gly Thr Ser Ile His Trp His Gly Leu His
 130 135 140

10 Gln Lys Gly Thr Asn Tyr His Asp Gly Ala Asn Gly Val Thr Glu Cys
 145 150 155 160

15 Pro Ile Pro Pro Gly Gly Ser Arg Val Tyr Ser Phe Arg Ala Arg Gln
 165 170 175

Tyr Gly Thr Ser Trp Tyr His Ser His Phe Ser Ala Gln Tyr Gly Asn
 180 185 190

20 Gly Val Ser Gly Ala Ile Gln Ile Asn Gly Pro Ala Ser Leu Pro Tyr
 195 200 205

Asp Ile Asp Leu Gly Val Leu Pro Leu Xaa Asp Trp Tyr Tyr Lys Ser
 210 215 220

25 Ala Asp Gln Leu Val Ile Glu Thr Leu Xaa Lys Gly Asn Ala Pro Phe
 225 230 235 240

30 Ser Asp Asn Val Leu Ile Asn Gly Thr Ala Lys His Pro Thr Thr Gly
 245 250 255

Glu Gly Glu Tyr Ala Ile Val Lys Leu Thr Pro Asp Lys Arg His Arg
 260 265 270

35 Leu Arg Leu Ile Asn Met Ser Val Glu Asn His Phe Gln Val Ser Leu
 275 280 285

Ala Lys His Thr Met Thr Val Ile Ala Ala Asp Met Val Pro Val Asn
 290 295 300

40 Ala Met Thr Val Asp Ser Leu Phe Met Ala Val Gly Gln Arg Tyr Asp
 305 310 315 320

45 Val Thr Ile Asp Ala Ser Gln Ala Val Gly Asn Tyr Trp Phe Asn Ile
 325 330 335

Thr Phe Gly Gly Gln Gln Lys Cys Gly Phe Ser His Asn Pro Ala Pro
 340 345 350

5 Ala Ala Ile Phe Arg Tyr Glu Gly Ala Pro Asp Ala Leu Pro Thr Asp
 355 360 365

Pro Gly Ala Ala Pro Lys Asp His Gln Cys Leu Asp Thr Leu Asp Leu
 370 375 380

10 Ser Pro Val Val Gln Lys Asn Val Pro Val Asp Gly Phe Val Lys Glu
 385 390 395 400

Pro Gly Asn Thr Leu Pro Val Thr Leu His Val Asp Gln Ala Ala Ala
 15 405 410 415

Pro His Val Phe Thr Trp Lys Ile Asn Gly Ser Ala Ala Asp Val Asp
 420 425 430

20 Trp Asp Arg Pro Val Leu Glu Tyr Val Met Asn Asn Asp Leu Ser Ser
 435 440 445

Ile Pro Val Lys Asn Asn Ile Val Arg Val Asp Gly Val Asn Glu Trp
 450 455 460

25 Thr Tyr Trp Leu Val Glu Asn Asp Pro Glu Gly Arg Leu Ser Leu Pro
 465 470 475 480

His Pro Met His Leu His Gly His Asp Phe Phe Val Leu Gly Arg Ser
 30 485 490 495

Pro Asp Val Ser Pro Asp Ser Glu Thr Arg Phe Val Phe Asp Pro Ala
 500 505 510

35 Val Asp Leu Pro Arg Leu Arg Gly His Asn Pro Val Arg Arg Asp Val
 515 520 525

Thr Met Leu Pro Ala Arg Gly Trp Leu Leu Leu Ala Phe Arg Thr Asp
 530 535 540

40 Asn Pro Gly Ala Trp Leu Phe His Cys His Ile Ala Xaa His Val Ser
 545 550 555 560

Gly Gly Leu Ser Val Asp Phe Leu Glu Arg Pro Asp Glu Leu Arg Gly
 45 565 570 575

Gln Leu Thr Gly Glu Ser Lys Ala Glu Leu Glu Arg Val Cys Arg Glu
580 585 590

5 Trp Lys Asp Trp Glu Ala Lys Ser Pro His Gly Lys Ile Asp Ser Gly
595 600 605

Leu Lys Gln Arg Arg Trp Asp Ala
610 615

10

CLAIMS

1. A variant of a parent laccase, which variant has laccase activity, improved stability as compared to said parent laccase, and comprises a mutation in one or more tyrosine, tryptophan or methionine residues, wherein the parent laccase has the amino acid sequence given in SEQ ID No. 1 or the parent laccase has an amino acid sequence which is at least 80% homologous to SEQ ID No. 1.
- 10 2. A variant according to claim 1, which comprises a mutation in a position corresponding to at least one of the following positions in SEQ ID No. 1:
M433,
15 W373,
W136,
Y145,
M480,
Y137,
20 Y176,
M254, and/or
W507.
3. A variant according to claim 1, wherein the parent laccase is derived from *Myceliophthora*.
- 25 4. A variant according to claim 1, wherein the parent laccase is derived from *Scytalidium*.
- 30 5. A variant according to claim 4, wherein the parent laccase is a *Scytalidium thermophilum* laccase with the sequence ID No. 2.
6. A variant according to claim 5, which comprises a mutation in a position corresponding to at least one of the following positions in SEQ ID No. 2:
M483,

W422,
W181,
Y190,
M530,
5 Y182,
Y221,
M300, and/or
M313.

10 7. A DNA construct comprising a DNA sequence encoding a laccase variant according to any of claims 1-6.

8. A recombinant expression vector which carries a DNA construct according to claim 7.

15

9. A cell which is transformed with a DNA construct according to claim 7 or a vector according to claim 8.

10. A cell according to claim 9, which is a microorganism.

20

11. A cell according to claim 10, which is a bacterium or a fungus.

12. A cell according to claim 11, which is an *Aspergillus niger*
25 or an *Aspergillus oryzae* cell.

13. Use of a laccase variant according to any of claims 1-6 for oxidizing a substrate.

30 14. Use of a laccase variant according to claim 13 for dye transfer inhibition.

15. Use of a laccase variant according to claim 13 for bleaching textiles, in particular for bleaching denim.

35

16. A detergent additive comprising a laccase variant according to any of claims 1-6 in the form of a non-dusting granulate, a stabilised liquid or a protected enzyme.

17. A detergent additive according to claim 16, which additionally comprises one or more other enzyme such as a protease, a lipase, an amylase, and/or a cellulase.

5

18. A detergent composition comprising a laccase variant according to any of claims 1-6 and a surfactant.

19. A detergent composition according to claim 18 which additionally comprises one or more other enzymes such as a protease, a lipase, an amylase and/or a cellulase.

10

THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 97/00566

A. CLASSIFICATION OF SUBJECT MATTER		
<p>IPC6: C12N 9/02 // (C12N 9/02, C12R 1:645) According to International Patent Classification (IPC) or to both national classification and IPC</p>		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
IPC6: C12N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
SE,DK,FI,NO classes as above		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9533836 A1 (NOVO NORDISK BIOTECH, INC. ET AL), 14 December 1995 (14.12.95)	1-19
	--	
X	WO 9533837 A1 (NOVO NORDISK BIOTECH, INC.), 14 December 1995 (14.12.95)	1-19
	--	

<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
<p>* Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>		
Date of the actual completion of the international search		Date of mailing of the international search report
30 March 1998		01 -04- 1998
Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86		Authorized officer Patrick Andersson Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

Information on patent family members

02/03/98

International application No.

PCT/DK 97/00566

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	9533836	A1	14/12/95	AU	2656595 A	04/01/96
				CA	2191718 A	14/12/95
				EP	0765394 A	02/04/97
				FI	964808 A	02/12/96
WO	9533837	A1	14/12/95	AU	2656695 A	04/01/96
				EP	0763115 A	19/03/97